

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF UTAH  
**Introduction to Mathematical Finance**  
**MATH 5760/6890 – Section 001 – Fall 2024**  
**Homework 11 Solutions**  
**Brownian Motion**

**Due: Friday, Nov 22, 2024**

---

Submit your homework assignment on Canvas via Gradescope.

- 1.) Recall that Brownian motion, which we identify with the log-return of an asset price, does not have bounded variation. Here is an exercise to motivate why this is consistent with our finance models: Let  $T, \mu, \sigma, S_0$  be fixed, with  $T, \sigma, S_0 > 0$ . Consider a real-world CRR model for the price  $S_n$  of the security  $S$ . Show that in the limit as  $n \rightarrow \infty$ , the *variation* of the log-returns of the real-world CRR trajectory is unbounded:

$$\lim_{n \rightarrow \infty} \sum_{j=1}^n |\log S_j - \log S_{j-1}| = \infty.$$

(This statement is true with probability 1.)

**Solution:** The CRR binomial tree model is that  $L_j = \log S_{j+1} - \log S_j$  is a Bernoulli( $p_n$ ) random variable taking values  $\log u_n$  with probability  $p_n$  and  $\log d_n$  with probability  $1 - p_n$ . The real-world CRR equations prescribe the values for  $(p_n, u_n, d_n)$ :

$$u_n = \exp(\sigma\sqrt{h_n}), \quad d_n = \exp(-\sigma\sqrt{h_n}), \quad p_n = \frac{1}{2} \left( 1 + \frac{\mu}{\sigma} \sqrt{h_n} \right),$$

where  $h_n = T/n$ . Note then that,

$$|\log S_j - \log S_{j-1}| = \left| \log \frac{S_j}{S_{j-1}} \right| = |\log L_{j-1}| = \begin{cases} |\log u_n|, & \text{with probability } p_n \\ |\log d_n|, & \text{with probability } 1 - p_n \end{cases}$$

However, the real-world CRR equations imply,

$$|\log u_n| = |\log d_n| = \sigma\sqrt{h_n}.$$

Hence, with probability 1,

$$|\log S_j - \log S_{j-1}| = \sigma\sqrt{h_n},$$

and therefore,

$$\begin{aligned} \lim_{n \rightarrow \infty} \sum_{j=1}^n |\log S_j - \log S_{j-1}| &= \lim_{n \rightarrow \infty} \sum_{j=1}^n \sigma\sqrt{h_n} = \sigma \lim_{n \rightarrow \infty} n\sqrt{h_n} \\ &= \sigma \lim_{n \rightarrow \infty} \sqrt{Tn} = \infty, \end{aligned}$$

as claimed.

- 2.) In this exercise, you will provide some evidence to support the fact that  $[B]_T = T$ , where  $B(t)$  is a standard Brownian motion. Define a discrete-time approximation of the quadratic variation of Brownian motion as,

$$Q_n := \sum_{j=1}^n (B(t_j) - B(t_{j-1}))^2, \quad t_j = jh_n, \quad h_n := \frac{T}{n}.$$

Show that  $Q_n$  has the following first- and second-order statistics for a fixed  $n$ :

$$\mathbb{E}Q_n = T, \quad \text{Var}Q_n = \frac{2T^2}{n}.$$

(Hence, as  $n \rightarrow \infty$ ,  $Q_n$  converges to mean- $T$  random variable with variance 0.)

Here is a helpful fact to aid computations: if  $X \sim \mathcal{N}(0, 1)$ , then  $\mathbb{E}X^4 = 3$ .

**Solution:** First, we note that by standard properties of Brownian motion:

$$B(t_j) - B(t_{j-1}) \sim \mathcal{N}(0, h_n),$$

and that

$$X_j := B(t_j) - B(t_{j-1}), \quad \{X_j\}_{j=1}^n \text{ are iid.}$$

We will utilize the hint at this time: since  $X_j \sim \mathcal{N}(0, h_n)$ , then  $\frac{1}{\sqrt{h_n}}X_j \sim \mathcal{N}(0, 1)$ , and so

$$\frac{1}{h_n^2} \mathbb{E}X_j^4 = \mathbb{E} \left( \frac{1}{\sqrt{h_n}} X_j \right)^4 = 3 \implies \mathbb{E}X_j^4 = 3h_n^2.$$

Returning to computing statistics of  $Q_n$ , we have:

$$\mathbb{E}Q_n = \sum_{j=1}^n \mathbb{E}(B(t_j) - B(t_{j-1}))^2 = \sum_{j=1}^n \text{Var}X_j = nh_n = T,$$

which shows the desired first-order statistic equality. The second equality requires us to compute,

$$\text{Var}Q_n = \sum_{j=1}^n \text{Var}(B(t_j) - B(t_{j-1}))^2 = \sum_{j=1}^n \text{Var}X_j^2.$$

Note that to compute the desired value, we need to compute  $\text{Var}X_j^2$ , where  $X_j$  is a standard normal random variable. This can be directly:

$$\text{Var}X_j^2 = \mathbb{E}(X_j^2 - \mathbb{E}X_j^2)^2 = \mathbb{E}X_j^4 - (\mathbb{E}X_j^2)^2 = \mathbb{E}X_j^4 - h_n^2 = 3h_n^2 - h_n^2 = 2h_n^2.$$

Putting this back in our equation for  $\text{Var}Q_n$ , yields,

$$\text{Var}Q_n = \sum_{j=1}^n \text{Var}X_j^2 = \sum_{j=1}^n 2h_n^2 = 2n \frac{T^2}{n^2} = \frac{2T^2}{n},$$

again as desired.